## Fate Report for Case # P-19-0140

#### **Fate Summary Statement**

Fate Summary P-19-0140 Statement: FATE: with MP < 25 °C (E)  $\log Kow = 4.50 (E)$  $S = 6.56 \text{ mg/L at } 25 \text{ }^{\circ}\text{C (E)}$ VP =torr at  $^{\circ}C$  (M) BP =°C (M) H = 1.79E-2 (E) $\log Koc = 2.67 (E)$  $\log Fish BCF = 2.64 (440) (E)$  $\log \text{ Fish BAF} = 3.38 (2,400) (E)$ POTW removal (%) = 50-90 via possible sorption and stripping; OECD 310(Headspace): 0.3%/28d NRB. Time for complete ultimate aerobic biodeg > mo Sorption to soils/sediments = low - moderate Volatilization half-life from a standard river = 2 hrs

Sorption to soils/sediments = low - moderate
Volatilization half-life from a standard river = 2 hrs
Volatilization half-life from a standard lake = 7 da
Atmospheric Oxidation Half-life = 97 hr via OH radical
Atmospheric Oxidation Half-life = 490 hr via ozone
PBT Potential: P3BU

FATE: Migration to ground water = moderate - rapid

Analogue found (include identifier and database)

Relevant Structure(s)
Water Parent
Landfill Parent
Air / Incineration Parent
Parent % incineration 99.9% (hazardous waste incinerator)

**Environmental Fate Determination** 

PMN #: P-19-0140

Summary: The new chemical substance is estimated to be very persistent in the environment and its bioaccumulation potential is unknown. Thus, it is unknown whether repeated exposures will cause food-chain effects via accumulation in exposed organisms.

Fate: Environmental fate is the determination of which environmental compartment(s) a chemical moves to, the expected residence time in the environmental compartment(s) and removal and degradation processes. Environmental fate is an important factor in determining exposure and thus in

determining whether a chemical may present an unreasonable risk. EPA estimated physical/chemical and fate properties of the new chemical substance using data for analogue(s) (highly fluorinated aliphatic chemicals), data submitted for the new chemical substance, and EPI (Estimation Program Interface) Suite<sup>TM</sup> (http://www.epa.gov/tsca-screening-tools/epi-suitetm-estimation-program-interface). In wastewater treatment, the new chemical substance is expected to be removed with an efficiency of 50% to 90% due to stripping and possible sorption. Removal of the new chemical substance by biodegradation is negligible. Sorption of the new chemical substance to sludge, soil, and sediment is expected to be low to moderate. Migration of the new chemical substance to groundwater is expected to be moderate to rapid due to low to moderate sorption to soil and sediment. Due to moderate reported vapor pressure, the new chemical substance is expected to undergo moderate volatilization to air. Overall, these estimates indicate that the new chemical substance has moderate potential to volatilize to air and has moderate to high potential to migrate to groundwater.

Persistence: Persistence is relevant to whether a new chemical substance is likely to present an unreasonable risk because chemicals that are not degraded in the environment at rates that prevent substantial buildup in the environment, and thus increase potential for exposure, may present a risk if the substance presents a hazard to human health or the environment. EPA estimated degradation half-lives of the new chemical substance using data for analogue(s) (highly fluorinated aliphatic chemicals) and data submitted for the new chemical substance. EPA estimated that the new chemical substance's aerobic and anaerobic biodegradation half-lives are > 6 months. These estimates indicate that the new chemical substance may be very persistent in aerobic environments (e.g., surface water) and anaerobic environments (e.g., sediment).

Bioaccumulation: Bioaccumulation is relevant to whether a new chemical substance is likely to present an unreasonable risk because substances that bioaccumulate in aquatic and/or terrestrial species pose the potential for elevated exposures to humans and other organisms via food chains. EPA estimated that the new chemical substance has unknown bioaccumulation potential based on uncertainty about bioaccumulation of the new chemical substance. The new chemical substance is estimated to be very persistent in the environment and its bioaccumulation potential is unknown. Thus, it is unknown whether repeated exposures will cause food-chain effects via accumulation in exposed organisms.

Fate Assessor: Card, Marcella
SMILES:

### **Physical Properties**

Property	Measured/Calculated Value	EPI
Molecular Form:	***	
Molecular Wt.:		g/mol
% < 500:		N957-

Property	Measured/Calculated Value	EPI
% < 1000:		

Property	Measured	Method	Estimated	Method	EPI
=======================================	Value		Value		
<b>Melting Point:</b>					NaN °C (Exp.)
					-65.33000183105469 °C (Est.,
					Joback)
					-71.07168579101562 °C (Est.,
					Gold)
					-68.20083618164062 °C (Est.,
D. III. D. I.	72.5				Selected)
<b>Boiling Point:</b>	73.5				NaN °C (Exp.)
					72.94 °C (Est.)
	<b>-</b> 60.0				346.1009216308594 °K (Est.)
<b>BP Pressure:</b>	760.0				
Vapor Pressure:	108.0				NaN mmHg (Exp.)
					17407.138431518553 Pa (Est.,
					Antoine)
					130.56463623046875 mmHg
					(Est., Antoine) 16572.619547720118 Pa (Est.,
					Grain)
					124.30521255096772 mmHg
					(Est., Grain)
					17718.11184888768 Pa (Est.,
					Mackay)
					132.89713512314304 mmHg
					(Est., Mackay)
					16989.878989619338 Pa (Est.,
					Selected)
					127.43492439071824 mmHg
					(Est., Selected)
					NaN Pa (Est., SubCooled)
					NaN mmHg (Est., SubCooled)
Water Solubility:			0.0066		NaN (Exp.)
					0.905242919921875 (Est.)
Log P:			4.5		
Log Kow:					NaN (Exp.)
					4.5 (Est.)
Log Koc:	NaN				6.153030553846979 (Est.,
					log(MCI))
					7.241731553111513 (Est.,
					log(Kow))
					470.1400146484375 L/kg (Est.,
					MCI)
					1396.510009765625 L/kg (Est.,
					Kow)
Log BCF:					3.16 L/kg wet-wt

Property	Measured Value	Method	Estimated Value	Method	EPI
Henry's Law:					NaN atm-m3/mole (Exp.)
					0.01789850927889347 atm-
					m3/mole (Est., Bond)
					0.0 atm-m3/mole (Est., Group)

TI.	
pH:	
II C	
pH Comment:	

# **Fate Analysis**

Hydrolysis (t1/2, da):		olatilization t1/2) - River		Volatilization (t1/2) - Lake
	`	(hr):		(da):
Atm Ox	8.076571794022694	Atm Ox	0.0560000017285347	Atm Ox
Potential		<b>Potential</b>		Potential(t1/2)
(t1/2)OH		(t1/2)O3		Total (hr):
(hr):		(hr):		
	0.10000000149011612	MITI	0.0	
Linear:		NonLinear:		
Biodeg		Biodeg	0.0	
Linear:		NonLinear:		
	1.2999999523162842	U	2.5999999046325684	
Survey ult:		Survey		
		Prim:		
	91.39601391219422%	,	0.18776511135574145%	
removal)		removal)		
Total:		Biodeg:		
	37.88274830106909%	,	53.325500940885746%	
removal)		removal)		
Ads:		Air:		

### Rationales

Removal in	
Wastewater	
<b>Treatment:</b>	
Atmospheric	
Oxidation:	
Hydrolysis:	
Photolysis:	
Aerobic	
<b>Biodegradation:</b>	
Anaerobic	

Biodegradation:
Sorption to Soil and
Sediment:
Migration to
Groundwater:
Persistence - Air:
Persistence Water:
Volatilization from
Water:
Soil:
Sediment:
Other:
Standard:
Bioaccumulation:

## **PBT Ratings**

Persistence	Bioaccumulation	Toxicity	PBT Comments
3	U	2 (HH, ECO)	

## **Exposure-Based Testing**

Exposure-Based	
Testing:	

### **Fate Ratings**

### Removal in WWT/POTW (Overall):

Removal in 50-90 WWT/POTW (Overall):

Condition	Rating Values		Rating Description			
		1	2	3	4	
WWT/POTW Sorption:	1-2	Low	Moderate	Strong	V. Strong	
WWT/POTW Stripping:	3	Extensive	Moderate	Low	Negligible	

Condition	Rating Values	Rating Description				Comment
		1	2	3	4	
Biodegradation Removal:	4	Unknown	High	Moderate	Negligible	
Biodegradation Destruction:		Unknown	Complete	Partial	_	
Aerobic Biodeg Ult:		<= Days	Weeks	Months	> Months	
Aerobic Biodeg Prim:		<= Days	Weeks	Months	> Months	
Anaerobic Biodeg Ult:		<= Days	Weeks	Months	> Months	
Anaerobic Biodeg Prim:		<= Days	Weeks	Months	> Months	
Hydrolysis (t1/2 at pH 7,25C) A:		<= Minutes	Hours	Days	>= Months	
Hydrolysis (t1/2 at pH 7,25C) B:		<= Minutes	Hours	Days	>= Months	
Sorption to Soils/Sediments:		V. Strong	Strong	Moderate	Low	
Migration to Ground Water:	3-4	Negligible	Slow	Moderate	Rapid	
Photolysis A, Direct:		Negligible	Slow	Moderate	Rapid	
Photolysis B, Indirect:		Negligible	Slow	Moderate	Rapid	
Atmospheric Ox A, OH:		Negligible	Slow	Moderate	Rapid	
Atmospheric Ox B, O3:		Negligible	Slow	Moderate	Rapid	

### **Bio Comments:**

**Bio Comments:** Fate Study Summaries are available.

Fish  $\log BAF = 3.38 (2,400)$ .

The fugacity spreadsheet and the EPI output file for the PMN material are attached.

### **Fate Comments:**

Fate Comments: OECD 310(Headspace): 0.3%/28d NRB.

### **Comments/Telephone Log**

A wtifa at	Undata/Unload Time
Artifact	Opdate/Opioad Time

Artifact	Update/Upload Time
De	